

9G8 Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP00050

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human SFRS7. AA range:61-110

Mol wt 27367

Species reactivity Human, Mouse

Clonality Polyclonal

Recommended application WB, IHC-p, ELISA

Concentration 1 mg/ml

Full name 9G8 Antibody

Synonyms SRSF7; SFRS7; Serine/arginine-rich splicing factor 7; Splicing factor 9G8; Splicing factor; arginine/serine-

rich 7

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


Serine and arginine rich splicing factor 7 encoded by SRSF7 is a member of the serine/arginine (SR)-rich family of pre-mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two transcript variants encoding different isoforms have been found for SRSF7.

Recommended Dilution

WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 ELISA: 1: 40000

Not yet tested in other applications.


Images




-- 26

Western Blot analysis of HuvEc cells using 9G8 Polyclonal Antibody cells nucleus.

Western blot analysis of lysates from HT-29 cells, using SFRS7 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using SFRS7 antibody.



Western Blot analysis of various cells using 9G8 Polyclonal Antibody cells nucleus.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com