

AID Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP00304
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human AICDA. AA range:81-130
Mol wt	23954
Species reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Recommended application	WB, IHC-p, ELISA
Concentration	1 mg/ml
Full name	AID Antibody
Synonyms	AICDA; AID; Activation-induced cytidine deaminase; Cytidine aminohydrolase

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

AICDA encodes a RNA-editing deaminase that is a member of the cytidine deaminase family. Activation-induced cytidine deaminase is involved in somatic hypermutation, gene conversion, and class-switch recombination of immunoglobulin genes. Defects in this gene are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2).

Recommended Dilution


WB: 1: 500 - 1: 2000

IHC-p: 1: 100 - 1: 300


ELISA: 1: 10000

Not yet tested in other applications.

Images



Western Blot analysis of HeLa, HepG2 cells using AID Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human colon cancer, antibody was diluted at 1:100

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com