

Histone deacetylase 6 Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP03986

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human HDAC6. AA range:1166-

1215

Mol wt 131431

Species reactivity Human, Mouse

Clonality Polyclonal

Recommended application WB, IHC-p, IF, ELISA

Concentration 1 mg/ml

Full name Histone deacetylase 6 Antibody

Synonyms HDAC6; KIAA0901; JM21; Histone deacetylase 6; HD6

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Histone deacetylase 6 encoded by HDAC6 belongs to class II of the histone deacetylase/acuc/apha family. It contains an internal duplication of two catalytic domains which appear to function independently of each other. This protein possesses histone deacetylase activity and represses transcription.

Recommended Dilution

WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 IF: 1: 200 - 1: 1000 ELISA: 1: 20000

Not yet tested in other applications.

Images




Western Blot analysis of MCF7 cells using Histone deacetylase 6 Polyclonal Antibody diluted at

1:2000



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using HDAC6 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using HDAC6 Antibody. The lane on the right is blocked with the synthesized peptide.

Western Blot analysis of various cells using Histone deacetylase 6 Polyclonal Antibody diluted at 1:2000

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com