

Histone H2A Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP03998

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human Histone H2A. AA range:1-50

Mol wt 51633

Species reactivity Human, Mouse, Rat

Clonality Polyclonal

Recommended application IHC-p, WB, ELISA

Concentration 1 mg/ml

Full name Histone H2A Antibody

Synonyms H2AFZ; H2AZ; Histone H2A.Z; H2A/z

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. H2AFZ encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic development and indicate that lack of functional histone H2A leads to embryonic lethality.

Recommended Dilution


WB: 1: 500 - 2000 ELISA: 1: 5000 IHC: 1: 100 - 1: 300

Not yet tested in other applications.

Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Histone H2A Antibody. The picture on the right is blocked with the synthesized peptide.

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com