

# PLC Gamma2 Polyclonal Antibody

### Description

Product type Primary Antibody

Code BT-AP07264

**Host** Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human PLCG2. AA range:721-770

Mol wt 147937

Species reactivity Human, Mouse, Rat

Clonality Polyclonal

Recommended application WB, IHC-p, ELISA

Concentration 1 mg/ml

Full name PLC gamma2 Antibody

Synonyms PLCG2; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesterase gamma-2; Phosphoinositide

phospholipase C-gamma-2; Phospholipase C-IV; PLC-IV; Phospholipase C-gamma-2; PLC-gamma-2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

### Background


The protein encoded by PLCG2 (phospholipase C gamma 2) is a transmembrane signaling enzyme that catalyzes the conversion of 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate to 1D-myo-inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG) using calcium as a cofactor. IP3 and DAG are second messenger molecules important for transmitting signals from growth factor receptors and immune system receptors across the cell membrane. Mutations in PLCG2 have been found in autoinflammation, antibody deficiency, and immune dysregulation syndrome and familial cold autoinflammatory syndrome 3.

## Recommended Dilution


WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 ELISA: 1: 5000

Not yet tested in other applications.

#### **Images**



Western blot analysis of lysates from COLO205 cells, using PLCG2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PLCG2 Antibody. The picture on the right is blocked with the synthesized peptide.



# Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com